

Marshall Islands

DEMOGRAPHICS







NUTRITIONAL STATUS

| Stunted (under-fives, 000): Underweight (under-fives, 000): .
| WHO Child Growth Standards | Stunted (under-fives, 000): Share of developing world stunting burden (%): Wasted (under-fives, 000): .
| Stunting country rank: Severely wasted (under-fives, 000): .

Current nutritional status

Percentage of children < 5 years old suffering

No Data

Stunting trends

Percentage of children < 5 years old stunted NCHS reference population

No Data

Underweight trends

Percentage of children < 5 years old underweight NCHS reference population


No Data

INFANT AND YOUNG CHILD FEEDING




Marshall Islands

MICRONUTRIENTS



lodized salt consumption trends*
Percentage of households consuming adequately iodized salt

No Data



ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE





MATERNAL NUTRITION/HEALTH

WATER AND SANITATION


Drinking water coverage

Under-five deaths caused by:





Diarrhoea:



DISPARITIES IN NUTRITION

Indicator	Gender			Residence			Wealth quintile						
	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-	-		-	-					-			
Underweight prevalence (WHO Child Growth Standards, %)	-		•	-				-		-		•	•
Wasting prevalence (WHO Child Growth Standards, %)		-	•	-	-	-				-	-	ě	•
Infants not weighed at birth (%)	14	16	0.9	8	28	0.3	32	25	10	9	8	0.3	DHS 2007
Early initiation of breastfeeding (%)	71	74	1.0	69	78	0.9	80	72	73	66	67	0.8	DHS 2007
Women with low BMI ($<$ 18.5 kg/m², %)				-	-					-			