

Country Profile

Nigeria

Maternal, Newborn & Child Survival

March 2012


Nigeria

DEMOGRAPHICS

Total population (000)	158,423	(2010)
Total under-five population (000)	26,569	(2010)
Births (000)	6,332	(2010)
Under-five mortality rate (per 1000 live births)	143	(2010)
Infant mortality rate (per 1000 live births)	88	(2010)
Neonatal mortality rate (per 1000 live births)	40	(2010)
Total under-five deaths (000)	861	(2010)
Maternal mortality ratio, adjusted (per 100,000 live births)	840	(2008)
Maternal mortality ratio, reported (per 100,000 live births)	545	(2002-2008)
Lifetime risk of maternal death (1 in N)	23	(2008)
Total maternal deaths (number)	50,000	(2008)

Under-five mortality rate


Deaths per 1000 live births




Causes of under-five deaths

Globally more than one third of child deaths are attributable to under nutrition

Causes of under-five deaths, 2008



Causes of neonatal deaths, 2008



INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION⁴

Wasting prevalence (based on 2006 WHO reference population, moderate and severe, %)	14	(2008)	Introduction of solid, semi-solid or soft foods (6-9 months, %)	76	(2008)	Low birthweight incidence (%)	12	(2008)
---	----	--------	---	----	--------	-------------------------------	----	--------

Underweight prevalence

Percent of children <5 years underweight for age


Based on 2006 WHO reference population



Stunting prevalence


Percent of children <5 years with low height for age

Based on 2006 WHO reference population



Exclusive breastfeeding


Percent of infants <6 months exclusively breastfed



CHILD HEALTH


Immunization

Percent of children immunised against measles
Percent of children immunised with 3 doses DPT
Percent of children immunised with 3 doses of Hib




Pneumonia treatment

Percent of children <5 years with suspected pneumonia taken to appropriate health provider
Percent of children <5 years with suspected pneumonia receiving antibiotics




Vitamin A supplementation

Percent of children 6-59 months receiving two doses of vitamin A during calendar year




Diarrhoeal disease treatment

Percent of children <5 years with diarrhoea receiving oral rehydration therapy (ORS, recommended homemade fluids or increased fluids), with continued feeding




Malaria treatment

Percent of febrile children <5 years using anti-malarials



Malaria prevention

Percent of children <5 years sleeping under ITNs




Nigeria

MATERNAL AND NEWBORN HEALTH


Proportion of women with low BMI (< 18.5 Kg/m2, %)	15	(2003)
Demand for family planning satisfied (%)	29	(2008)
Total fertility rate	5.5	(2010)
Adolescent birth rate (births per 1000 woman aged 15-19 yr)	123	(2003-)
Antenatal visit for woman (4 or more visits, %)	45	(2008)
Early initiation of breastfeeding (within 1 hour of birth, %)	38	(2008)
Institutional deliveries (%)	35	(2008)
Postnatal visit for baby (within 2 days for home births, %)	-	-
Postnatal visit for mother (within 2 days, %)	38	(2008)

Causes of maternal deaths
Regional estimates for sub-Saharan Africa, 1997-2007



Source: WHO 2010 Note: Figures may not add to 100% due to rounding.


Coverage along the continuum of care



Source: DHS, MICS, Other NS


Antenatal care

Percent of women aged 15-49 years attended at least once by a skilled health provider during pregnancy




Skilled attendant at delivery

Percent of live births attended by skilled health personnel



Neonatal tetanus protection

Percent of newborns protected against tetanus




Source: WHO/UNICEF 2011

HIV AND AIDS

HIV prevalence among young women (15-24 yrs, %)	2.9 [2.3 - 3.9]	(2009)
HIV prevalence among young men (15-24 yrs, %)	1.2 [0.9 - 1.6]	(2009)
HIV+ children receiving ART (%)	10 [7 - 19]	(2009)
Orphan school attendance ratio	1.17	(2008)

Prevention of mother-to-child transmission of HIV
Percent of HIV+ pregnant women receiving ARVs for PMTCT



Source: MOH/UNAIDS/WHO


EDUCATION

Survival to last grade of primary school (total, admin data, %)	-	-
Survival to last grade of primary school (male, admin data, %)	-	-
Survival to last grade of primary school (female, admin data, %)	-	-
Primary school net enrolment ratio (total, admin data, %)	63	(2007)
Primary school net enrolment ratio (male, admin data, %)	66	(2007)
Primary school net enrolment ratio (female, admin data, %)	60	(2007)

WATER AND SANITATION

Drinking water coverage


Percent of population by type of drinking water source, 2010



Source: WHO/UNICEF JMP 2012

Sanitation coverage

Percent of population by type of sanitation facility, 2010



Source: WHO/UNICEF JMP 2012

CHILD PROTECTION

Women aged 20-24 years who were married or in union by age 18 (%)	39	(2008)
Birth registration (%)	30	(2008)
Female genital mutilation/cutting (%)	30	(2008)

DISPARITIES IN INTERVENTION COVERAGE ¹

Indicator	Total	Gender			Residence			Wealth Quintile						Ratio of Richest to Poorest	Equity chart ²	Source
		Male	Female	Ratio of Male to Female	Urban	Rural	Ratio of Urban to Rural	Poorest	Second	Middle	Fourth	Richest				
DEMOGRAPHICS ³																
Under-five mortality rate (per 1,000 live births)	143	-	-	-	-	-	-	219	212	165	129	87	0.4		IGME 2011	
NUTRITION ⁴																
Low birth weight incidence (%)	12	-	-	-	11	12	0.9	14	12	11	10	10	0.7		DHS 2008	
Underweight prevalence (based on 2006 WHO reference population, %)	27	25	22	1.1	16	27	0.6	35	29	22	17	10	0.3		DHS 2008	
Stunting prevalence (based on 2006 WHO reference population, %)	41	43	38	1.1	31	45	0.7	52	49	42	34	24	0.5		DHS 2008	
Wasting prevalence (based on 2006 WHO reference population, %)	14	14	13	1.1	11	15	0.7	21	17	12	10	9	0.5		DHS 2008	
Exclusive breastfeeding (0-5 months, %)	13	12	14	0.9	19	11	1.8	7	8	12	20	21	3.1		DHS 2008	
Introduction of solid, semi-solid or soft foods (6-9 months, %)	76	75	78	1.0	82	74	1.1	74	71	75	83	83	1.1		DHS 2008	
Household consuming adequately iodized salt (15 ppm or more, %)	97	-	-	-	99	97	1.0	-	-	-	-	-	-		DHS 2003	
CHILD HEALTH ⁵																
Care seeking for pneumonia (%)	45	44	47	0.9	46	45	1.0	32	40	56	61	-	-		DHS 2008	
Antibiotic use for pneumonia (%)	23	24	21	1.1	24	22	1.1	12	21	28	33	-	-		DHS 2008	
Diarrhoeal treatment - children receiving ORT and continued feeding (%)	25	25	25	1.0	34	22	1.6	17	21	31	34	41	2.5		DHS 2008	
Malaria prevention - children sleeping under ITNs (%)	29	-	-	-	23	31	0.7	-	-	-	-	-	-		Other NS 2010	
Malaria treatment - febrile children receiving antimalarial medicines (%)	49	-	-	-	60	46	1.3	-	-	-	-	-	-		Other NS 2010	
MATERNAL AND NEWBORN HEALTH																
Antenatal care coverage at least one visit (%)	58	-	-	-	84	46	1.8	24	40	64	82	94	3.9		DHS 2008	
Antenatal care coverage (4 or more visits, %)	45	-	-	-	69	34	2.0	-	-	-	-	-	-		DHS 2008	
Skilled attendant at delivery (%)	39	-	-	-	65	28	2.4	8	18	38	63	86	10.3		DHS 2008	
Early initiation of breastfeeding (%)	38	38	39	1.0	41	38	1.1	30	38	43	43	41	1.4		DHS 2008	
WATER AND SANITATION ⁶																
Use of improved drinking water sources (%)	58	-	-	-	74	43	1.7	24	37	53	76	90	3.8		2010 (WHO/UNICEF JMP 2012)	
Use of improved sanitation facilities (%)	31	-	-	-	35	27	1.3	25	38	44	67	92	3.8		2010 (WHO/UNICEF JMP 2012)	
EDUCATION																
Survival rate to last grade of primary school (survey data, %)	98	98	98	1.0	98	-	-	-	-	-	-	-	-		DHS 2008	
Primary school net attendance ratio (survey data, %)	63	65	60	1.1	78	56	1.4	31	52	52	81	72	2.4		DHS 2008	
CHILD PROTECTION																
Women aged 20-24 years who were married or in union by age 18 (%)	39	-	-	-	22	50	0.4	71	65	43	23	11	0.1		DHS 2008	
Birth registration (%)	30	30	31	1.0	49	22	2.2	9	18	27	44	62	7.0		DHS 2008	
Female genital mutilation/cutting (%)	30	-	-	-	37	26	1.4	13	23	30	39	39	2.9		DHS 2008	

Note: The format for this Country Profile has been adapted from the Countdown to 2015 report. Coverage data have been largely derived from national household surveys such as the Multiple Indicator Cluster Surveys (MICS) and Demographic and Health Surveys (DHS). For the majority of coverage indicators, UNICEF global databases were used. Other organizations such as the World Health Organization, UNAIDS, United Nations Population Fund, London School of Hygiene and Tropical Medicine and Saving Newborn Lives also provided data. Details on indicators, data sources, and definitions of indicators, can be found at www.childinfo.org.

1. **Disparities** - Disparity information is only available for data directly derived from household surveys such as MICS and DHS. Therefore, disparity data are not available for the following indicators: vitamin A supplementation, immunization, and for HIV/AIDS. In addition, neither UNICEF Global Databases nor databases from partner organizations maintain disparity data for the following indicators: total fertility rate, unmet need, institutional deliveries, contraceptive prevalence, adolescent birth rate.

2. **Equity chart** - Displays values for the five wealth quintiles to the left. The scale is 0 to 100% for all charts except U5MR, which shows a range of 0 to 300 deaths per 1,000 live births.

3. **U5MR** - Wealth quintile data are derived directly from MICS, DHS or other surveys. The total is the inter-agency estimate published by the UN Inter-agency Group for Child Mortality Estimation (IGME).

4. **Anthropometric indicators - Reference Standards for Underweight, Stunting and Wasting.** New international Child Growth Standards for infants and young children were released by WHO in 2006, replacing the older NCHS/WHO reference population. In using the 2006 WHO reference population, estimates generally change in the following manner: stunting is greater throughout childhood; underweight rates are higher during the first half of infancy and lower thereafter; and, wasting rates are higher during infancy. Please note that there may be small discrepancies between the totals and the disparity data, as the totals have undergone additional analysis.

5. **Child Health** - All indicators in this section refer to children under 5 years of age.

6. **Water and sanitation** - Wealth quintile data are derived from MICS or DHS surveys. Urban, rural and total coverage estimates provided are for 2010 and are those published by the WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation.