



Countdown to 2015

Maternal, Newborn & Child Survival

Kyrgyzstan


May 2013

DEMOGRAPHICS

Total population (000)	5,393	(2011)
Total under-five population (000)	626	(2011)
Births (000)	131	(2011)
Birth registration (%)	94	(2009)
Total under-five deaths (000)	4	(2011)
Neonatal deaths: % of all under-5 deaths	48	(2011)
Neonatal mortality rate (per 1000 live births)	16	(2011)
Infant mortality rate (per 1000 live births)	27	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	91	(2010)
Lifetime risk of maternal death (1 in N)	480	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	31	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (moderate and severe, %)


Early initiation of breastfeeding (within 1 hr of birth, %)

Introduction of solid, semi-solid/soft foods (%)

Vitamin A two dose coverage (%)

Percent children <5 years who are underweight

Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

