

Central Africa

WUENIC 2025 revision,
Published 15 July 2026



WHO/UNICEF Estimates of National Immunization Coverage (WUENIC), 2025 revision

Every year, WHO and UNICEF jointly review submissions from Member States on national immunization coverage, including annual administrative and official coverage, finalized survey reports and data from both published and grey literature. The data is triangulated with consideration of potential biases and local expert opinions to differentiate between accurately reflective empirical data and potentially misleading data, to assess the most likely coverage levels for each country.

WHO and UNICEF produce country-specific estimates by reviewing each country's data independently, without borrowing from other countries when data are missing. The estimates are not ad hoc adjustments to reported figures and may rely on a single empirical source, typically nationally reported coverage. When no data exist for a specific country-vaccine-year, earlier and later data are interpolated. If sources conflict or vary widely, the most plausible estimate is selected after considering potential biases.

This slide deck presents the latest WUENIC estimates (published 15 July 2026), using UNPD World Population Prospects (WPP 2024) for population-weighted averages and to derive absolute numbers of vaccinated and unvaccinated/zero-dose children.

Methods: • [Burton et al. 2009. WHO and UNICEF estimates of national infant immunization coverage: methods and processes.](#)

- [Burton et al. 2012. A formal representation of the WHO and UNICEF estimates of national immunization coverage: a computational logic approach.](#)
- [Brown et al. 2013. An introduction to the grade of confidence used to characterize uncertainty around the WHO and UNICEF estimates of national immunization coverage.](#)
- [Danovaro-Holliday et al. 2021. Compliance of WUENIC with Guidelines for Accurate and Transparent Health Estimates Reporting \(GATHER\) criteria.](#)

Definitions of immunization terms

Vaccine coverage

Percentage of infants (children under one year of age) who received certain vaccine-doses. For example, coverage of DTP3 is the percentage of infants who received all three doses of diphtheria, tetanus, and pertussis (DTP) vaccine.

Unvaccinated

An infant that did not receive the first dose of a vaccine series. The term zero-dose is used to describe children unvaccinated with DTP1.

Under-vaccinated

An infant who received some but not all the recommended vaccine-doses in the national schedule.

Vaccine-Doses

- Bacillus Calmette-Guerin (BCG): vaccine against tuberculosis
- Hepatitis B birth dose, given within 24 hours after birth (HepBB)
- Diphtheria, tetanus, and pertussis vaccine, first dose (DTP1) and third dose (DTP3)
- Hepatitis B vaccine, third dose (HepB3)
- Haemophilus influenzae type b vaccine, third dose (Hib3)
- Inactivated polio vaccine, first dose (IPV1) and last dose (IPVC)
- Measles containing vaccine, first dose (MCV1) and second dose (MCV2)
- Rotavirus vaccine, last dose (RotaC)
- Pneumococcal vaccine, last dose (PCVC)
- Yellow Fever vaccine (YFV)
- Meningococcal A vaccine (MengA)
- Human papillomavirus vaccine, first dose (HPV1) and last dose (HPVc): vaccine to protect against certain types of human papillomavirus that can lead to cancer or genital warts

The Immunization Agenda 2030 (IA2030)

The IA2030 is a global strategy endorsed by the World Health Assembly aiming to ensure everyone, everywhere, at every age benefits from vaccines for improved health and well-being by 2030. It focuses on increasing vaccine coverage, equity, sustainability and pandemic preparedness while promoting life-course immunization and integrating immunization with other health services.

Key concepts


- The World Health Organization (WHO) provides global vaccine recommendations, which are adapted by countries based on local needs. Only DTP, polio and measles-containing vaccines are used in all countries.
- DTP1 is a marker of access to routine immunization services, and when not received, serves as a proxy for identifying children who have not received any vaccinations, also known as "zero-dose" children. High DTP1 coverage indicates good access to immunization services, while low coverage suggests challenges in reaching children with essential vaccines.
- DTP3 is a widely used indicator of immunization programme performance. It reflects a country's ability to deliver routine immunization services and ensures children are protected against serious disease. DTP3 is tracked globally and serves as a key measure of a nation's vaccination efforts.
- DTP1-DTP3 drop-out measures the percentage of children who received DTP1 but not DTP3, and highlights where children are lost along the vaccination pathway, highlighting potential weaknesses in service delivery and follow-up.
- MCV1 (usually recommended between 9-12 months) assesses the ability to deliver vaccines later in infancy. It serves as a tracer for protection against measles and is a good indicator of health system performance.
- HPV vaccine protects against specific types of human papilloma virus (HPV), and is used to measure life cycle vaccination.
- Other key indicators include PCVC and MCV2, which are used to monitor the Sustainable Development Goals (SDGs).
- Together, these indicators provide a consistent and comparable way to track immunization progress, identify missed communities and monitor global targets, including those under the Immunization Agenda 2030 (IA2030) and Sustainable Development Goals (SDGs).

Overview

- DTP1 coverage increased 1 percentage point from 82% in 2024 to 83% in 2025.
- DTP3 coverage increased 2 percentage points from 68% in 2024 to 70% in 2025.
- In Central Africa, there were 68,000 fewer zero-dose children compared to in 2024. This leaves 1.2 million children without vaccination, vulnerable to vaccine-preventable diseases and a further 954,000 with only partial protection.
- In Central Africa, 1 countries* accounted for over half (63.8%) of all zero-dose children.
- There were 0 (0%) countries that achieved DTP3 coverage of at least 90% (IA2030 target).
- MCV1 coverage increased 1 percentage point from 61% in 2024 to 62% in 2025. There were 2.8 million children who missed out on the first measles vaccination.
- MCV2 coverage increased 3 percentage points from 50% in 2024 to 53% in 2025.
- Last dose coverage of HPV vaccination (HPVc) among girls decreased from 6% to 5% in 2025.

*DRC (63.8%)

Vaccine coverage (%), by year, Central Africa, 2000-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Vaccine coverage and progress varies between antigens and over time.


This heatmap shows vaccine coverage trends in Central Africa since 2000, with green cells indicating coverage of 90% or more.

In 2015, none of the 14 (0%) vaccines with data achieved coverage of 90% or more.

In 2025, none of the 16 (0%) vaccines achieved coverage of 90% or more.

Since 2006, estimates have been made for 6 new vaccines. IPV1 is the newest vaccine being reported (2021), which achieved 54% coverage in 2025.

Vaccine coverage, by country, Central Africa, 2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Countries ordered based on descending crude average coverage across all vaccines.

This heatmap shows vaccine coverage by country and vaccine in 2025.


In 2025, 4 out of 9 countries achieved coverage of 90% or higher in at least one vaccine.

Chad, Congo, and Sao Tome and Principe had the highest percentage of vaccines (8%) in the national immunization schedule with coverage of at least 90%, followed by Cameroon (7%).

5 countries had not achieved 90% coverage of any vaccine (Burundi, CAR, DRC, Equatorial Guinea, and Gabon).

DTP1

In 2025, DTP1 coverage was 83% and DTP3 was 70%, leaving 2.2 million children un- or under-vaccinated, including 1.2 million zero-dose and 954,000 with only partial protection.



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: There may be discrepancies due to rounding.


The key goal of the Immunization Agenda 2030 is to make vaccination available to everyone, everywhere, by 2030.

This chart shows diphtheria, tetanus and pertussis-containing vaccine first (DTP1) and third dose (DTP3) coverage trends, the number of zero-dose children and DTP drop-out.

In 2025, DTP1 coverage remained relatively constant within 1% at 83%. The number of children missing out on any vaccination (zero-dose children) improved from 1,303,000 in 2024 to 1,236,000 in 2025.

DTP3 coverage increased at 70% in 2025, leaving 2,190,000 children vulnerable to vaccine-preventable diseases.

In 2025, DTP1 coverage was 83%: 6 million children received at least one dose of DTP-containing vaccine, but 1.2 million remain unreached.



Sources: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision; United Nations, Department of Economic and Social Affairs, Population Division (2024). World Population Prospects 2024, Online Edition.

DTP1 coverage in 2025 (83%) was higher than in 2019 (80%).

The number of children vaccinated with DTP1 increased 18% from 5.1 million in 2019 to 6 million in 2025.


The number of surviving infants increased approximately 14% from 6.3 million in 2019 to 7.2 million in 2025.

In 2025, 911,000 more children were vaccinated than in 2019.

In 2025, there were 907,000 more surviving infants (target population) than in 2019.

For vaccine coverage to increase, the number of children vaccinated must increase at a faster rate than the population increases.

Countries ranked by number of zero-dose children, Central Africa, 2021-2025



This chart compares the ranking of countries in Central Africa based on the absolute number of zero-dose children, with rank 1 representing the country in the region with the most zero-dose children.


In 2021, DRC had the highest number of zero-dose children (757,000), followed by Cameroon (216,000).

In 2025, DRC had the highest number of zero-dose children (788,000), followed by Cameroon (132,000).

There was some change in the top 3 countries between 2021 and 2025.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Number in parentheses is the number of zero-dose children.
Zero-dose children are those who did not receive DTP1.

DTP coverage (%), number of children fully, partially and unvaccinated for DTP 2000-2025 and projections to 2030 based on IA2030 target , Central Africa




IA2030 calls on all countries to reduce the number of zero dose children in 2019 by half by 2030. This chart shows the annual number of children required to be vaccinated to reach the ZD target.

Central Africa is projected to have a moderate increase (400,000-1 million) in the number of surviving infants by 2030. Therefore, maintaining current coverage requires vaccinating an increasing number of children.

To achieve the IA2030 ZD target, more (~3.68%) children need to be vaccinated with DTP1 each year. This will require increases in immunization programme and health system capacity.

Sources: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision; United Nations, Department of Economic and Social Affairs, Population Division (2024), World Population Prospects 2024, Online Edition. Note: The Immunization Agenda 2030 (IA2030) calls on all countries to reduce the number of zero dose children in 2019 by half by 2030.

In 2025, the number of zero-dose children was 33% above the annual IA2030 milestone, putting the 2030 goal at serious risk without accelerated action.




The Immunization Agenda 2030 (IA2030) aims to leave no one behind with immunization and calls on all countries to reduce the number of zero dose children by half by 2030.

- This chart shows:
- Estimated number of zero-dose children in 2000-2024 (dark blue bars)
 - Zero-dose target by 2030 (light blue bar)
 - Trajectory to reach the 2030 target based on a linear decline (points)

In 2025, the number of zero-dose children was approximately 33% higher than the annual milestone to reach the target.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: The Immunization Agenda 2030 (IA2030) calls on all countries to reduce the number of zero dose children in 2019 by half by 2030. The line and points show the yearly progress and trajectory to meet the target by 2030, based on a linear decline.

DTP1 coverage, by country, Central Africa, 2000-2025




Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Countries ordered based on descending coverage in 2025


This heatmap shows DTP1 coverage, by region and year. Red indicates low coverage (<60%) and green indicates high coverage (≥90%). The changing colour gradient can be used to tracked changes in coverage over time and recovery from pandemic-related disruptions where applicable.

DTP1 coverage has improved since 2000 in all regions and exceeded pre-pandemic levels in over half of regions, but progress is uneven and gaps in zero-dose children persist.

DTP1 coverage, Central Africa, 2000-2025




DTP1 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of zero-dose children, Central Africa



In 2025, Central Africa ranked number 3 out of 5 AU regions, with DTP1 coverage of 83%.

DTP1 coverage (83%) was 3 percentage points higher than in 2019 (80%).

This equates to 1.2 million zero-dose children in 2025 compared to 1.2 million zero-dose children in 2019.


Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019.
There may be discrepancies in total Zero-dose children (millions) due to rounding.

DTP1 coverage improved between 2024 and 2025 in over half of regions and exceeded pre-pandemic levels in over half of regions, but progress is uneven and gaps in zero-dose children persist.

DTP1 coverage, Central Africa, 2018-2025




DTP1 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of zero-dose children, Central Africa




In 2025, Central Africa ranked number 3 out of 5 AU regions, with DTP1 coverage of 83%.

DTP1 coverage (83%) was 3 percentage points higher than in 2019 (80%).

This equates to 1.2 million zero-dose children in 2025 compared to 1.2 million zero-dose children in 2019.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019.
There may be discrepancies in total Zero-dose children (millions) due to rounding.

In 2025, 1 countries achieved at least 90% DTP1 coverage, while 2 had coverage below 70%.



DTP1 coverage (%) ■ <60 ■ 60-69 ■ 80-89 ■ 90-94

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision


Note: This map is stylized and based on an approximate scale. This map does not reflect a position by UNICEF or WHO on the legal status of any country or territory or the delimitation of any frontiers.

In Central Africa, 2 out of 9 (22%) countries had DTP1 coverage below 70% in 2025. These are CAR and Gabon.

These countries account for approximately 305,000 (4%) of the total annual cohort of surviving infants in the region.

There were 1 (11%) countries that achieved DTP1 of 90% or higher, accounting for 191,000 (3%) of the total cohort.

Coverage of DTP1, by country, Central Africa, 2018-2025



In 2025, Congo had the highest coverage (93%), while CAR had the lowest coverage (59%).


DTP1 coverage increased in 6 out of 9 countries between 2024 and 2025. The largest increase was 9% points in Equatorial Guinea from 75% to 84%.

No countries saw a decrease in coverage. Coverage remained constant between 2024 and 2025 in the remaining 3 countries.

In 2025, 1 out of 9 countries had achieved DTP1 coverage of at least 90%.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Dark grey line displays regional coverage for reference

DTP1 coverage difference compared to 2019, by country, Central Africa, 2018-2025



This chart shows difference in DTP1 coverage compared to 2019, by country.

Coverage was higher in 2025 compared to 2019 in 5 countries (CAR, Cameroon, Chad, Congo, and Equatorial Guinea).

The largest increase in coverage was 17% points in Chad from 69% in 2019 to 86% in 2025.

Coverage was lower than in 2019 in 4 countries (Burundi, DRC, Gabon, and Sao Tome and Principe).


The largest decline in coverage was 12% points in Gabon from 77% in 2019 to 65% in 2025.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: Dark grey line displays regional coverage difference for reference

ence compared to 2019, values above zero indicate coverage n percentage points higher than in 2019 and values below zero indicate coverage n percentage points lower than in 2019

DTP1 coverage, by country, Central Africa, 2019-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: The grey bar spans vaccine coverage across all years 2019-2025 and the dots represent coverage in specific years. Countries are arranged by high to low coverage in 2025 and coloured based on if coverage is lower (red), the same as (blue) or higher (green) than in 2019. An asterisk (*) indicates countries aiming to catch-up children missed since 2019 through a Big Catch-up Plan, supported by financing from The Gavi Alliance.


This chart shows the range of DTP1 coverage across years 2019 to 2025 (grey bars), and coverage in specific years (dots), by country. The chart can be used for assessing recovery to pre-pandemic levels and progress beyond 2019.

In 2024, 5 (out of 9) countries had lower coverage than in 2019.

In 2025, 0 countries had lower DTP1 coverage than in 2024.

In 2025, 4 countries had lower coverage than in 2019.

Between 2019 and 2025, DTP1 coverage increased in five countries and decreased in four countries.




This chart shows DTP1 coverage in 2019 and 2025, and change in coverage between the two years.

Overall, 4 out of 9 (44%) countries in Central Africa had a decline in coverage. The largest decline was in Gabon (-12 percentage points), followed by Sao Tome and Principe (-10 percentage points).

There were 5 (56%) countries with an increase in DTP1 coverage. Coverage remained the same in 0 countries between 2019 and 2025.

Between 2024 and 2025, DTP1 coverage increased in six countries, and no countries experienced a decline.




This chart shows DTP1 coverage in 2024 and 2025, and change in coverage between the two years.

Overall, 0 out of 9 (0%) countries in Central Africa had a decline in coverage.

There were 6 (67%) countries with an increase in DTP1 coverage. Coverage remained the same in 3 countries between 2024 and 2025.

In 2025, CAR and Gabon had the lowest DTP1 coverage, while DRC and Cameroon had the highest number of zero-dose children, highlighting the need to consider both coverage and absolute numbers to identify all vulnerable populations.



This chart shows DTP1 coverage in countries in Central Africa from lowest to highest coverage, and the rank of the top 10 countries with the most zero-dose children, based on absolute numbers.

In 2025, DTP1 coverage ranged from 59% in CAR to 93% in Congo.

1 out of 9 (11%) countries had coverage of at least 90%.


DRC had DTP1 coverage of 82% and the highest absolute number of zero-dose children in the region, followed by Cameroon.

Large cohort countries may have high numbers of zero-dose children despite high vaccine coverage. It is advantageous to look at both coverage and absolute numbers of unvaccinated children to ensure vulnerable countries with small birth cohorts are not missed.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Bars are ranked by ascending coverage.
Numbers in bubbles display top 10 rank based on absolute number of zero-dose children. Number in parentheses shows number of zero-dose children.

In 2025, 1 countries accounted for over half of all zero-dose children in Central Africa, with DRC contributing the highest share.

DTP1 unvaccinated (% of total unvaccinated)




In 2025, 1 countries accounted for over half (63.8%) of all zero-dose children in Central Africa.

DRC had the highest number of children who did not receive DTP1 (n=788,000), which is approximately 63.8% of all zero-dose children in Central Africa (n=1.2 million).

The number of unvaccinated children is influenced by both the birth cohort size and programme performance. As a result, countries with large birth cohorts may still have a high number of unvaccinated children, even if coverage rates are high. At the same time, countries with high coverage will also have a greater absolute number of children who are vaccinated. Please refer to the heatmaps for country-specific coverage.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Number in legend corresponds to region ranking based on highest to lowest absolute number of unvaccinated children and reflects order of pie slices.

Approximately 75% of zero-dose children in Central Africa live in just 3 countries, highlighting where intensified efforts could have the greatest regional impact on zero-dose numbers.




Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

This chart shows the number of zero-dose children in 2025, by country in Central Africa. Bars are coloured based on DTP1 coverage.

In 2025, DRC had the highest absolute number of zero-dose children, with 82% DTP1 coverage.

The number of unvaccinated children is influenced by both the birth cohort size and programme performance. As a result, countries with large birth cohorts may still have a high number of unvaccinated children, even if coverage rates are high. At the same time, countries with high coverage will also have a greater absolute number of children who are vaccinated. Please refer to the heatmaps for country-specific coverage.

Country trends in zero-dose children were heterogeneous, reinforcing the need to target resources and strategies to countries where the burden is increasing while sustaining gains elsewhere.



This chart shows the absolute change in number of zero-dose children, by country.


Overall, 3 countries had an increase in zero-dose children and 6 countries saw the number of zero-dose children decline between 2024 and 2025.

The three countries with the largest increase were: DRC (772,000 $\Rightarrow$ 788,000 [+16,000]), Burundi (50,000 $\Rightarrow$ 50,000 [+<500]), Sao Tome and Principe (<1,000 $\Rightarrow$ <1,000 [+<100]).

Cameroon had the largest decline in the number of zero-dose children from 168,000 in 2024 to 132,000 in 2025 (-36,000).

DTP3

In 2025, DTP3 coverage was 70%: 5 million children received three doses of DTP-containing vaccine, but 2.2 million remain un- or under-vaccinated.



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
United Nations, Department of Economic and Social Affairs, Population Division (2024). World Population Prospects 2024, Online Edition.
Note: Unvaccinated includes zero-dose and undervaccinated children.

DTP3 coverage in 2025 (70%) was the same as in 2019 (70%).

The number of children vaccinated with DTP3 increased 13% from 4.4 million in 2019 to 5 million in 2025.


The number of surviving infants increased approximately 14% from 6.3 million in 2019 to 7.2 million in 2025.

In 2025, 588,000 more children were vaccinated than in 2019.

In 2025, there were 907,000 more surviving infants (target population) than in 2019.

For vaccine coverage to increase, the number of children vaccinated must increase at a faster rate than the population increases.

DTP3 coverage, by country, Central Africa, 2000-2025




This heatmap shows DTP1 coverage, by region and year. Red indicates low coverage (<60%) and green indicates high coverage (≥90%). The changing colour gradient can be used to tracked changes in coverage over time and recovery from pandemic-related disruptions where applicable.


Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Countries ordered based on descending coverage in 2025

DTP3 coverage has improved since 2000 in all regions and exceeded pre-pandemic levels in over half of regions, but progress is uneven and gaps in undervaccinated children persist.

DTP3 coverage, Central Africa, 2000-2025




DTP3 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of un- and undervaccinated children, Central Africa



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019. There may be discrepancies in total Undervaccinated children (millions) due to rounding.


In 2025, Central Africa ranked number 5 out of 5 AU regions, with DTP3 coverage of 70%.

DTP3 coverage (70%) was the same as in 2019 (70%).


This equates to 2.2 million undervaccinated children in 2025 compared to 1.9 million undervaccinated children in 2019.

DTP3 coverage improved between 2024 and 2025 in over half of regions and exceeded pre-pandemic levels in over half of regions, but progress is uneven and gaps in undervaccinated children persist.

DTP3 coverage, Central Africa, 2018-2025




DTP3 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of un- and undervaccinated children, Central Africa



In 2025, Central Africa ranked number 5 out of 5 AU regions, with DTP3 coverage of 70%.


DTP3 coverage (70%) was the same as in 2019 (70%).

This equates to 2.2 million undervaccinated children in 2025 compared to 1.9 million undervaccinated children in 2019.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019.
There may be discrepancies in total Undervaccinated children (millions) due to rounding.

In 2025, 0 countries achieved at least 90% DTP3 coverage, while 3 had coverage below 70%.

DTP3 Coverage (%), Central Africa, 2025



DTP3 coverage (%) ■ <60 ■ 60-69 ■ 70-79 ■ 80-89

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision


Note: This map is stylized and based on an approximate scale. This map does not reflect a position by UNICEF or WHO on the legal status of any country or territory or the delimitation of any frontiers.

In Central Africa, 3 out of 9 (33%) countries had DTP3 coverage below 70% in 2025. These are CAR, DRC, and Gabon.

These countries account for approximately 4.7 million (65%) of the total annual cohort of surviving infants in the region.

There were 0 (0%) countries that achieved DTP3 of 90% or higher, accounting for <100 (0%) of the total cohort.

Coverage of DTP3, by country, Central Africa, 2018-2025




In 2025, Burundi had the highest coverage (89%), while CAR had the lowest coverage (46%).

DTP3 coverage increased in 6 out of 9 countries between 2024 and 2025. The largest increase was 10% points in Equatorial Guinea from 68% to 78%.

No countries saw a decrease in coverage. Coverage remained constant between 2024 and 2025 in the remaining 3 countries.

In 2025, 0 out of 9 countries had achieved DTP3 coverage of at least 90%.

DTP3 coverage difference compared to 2019, by country, Central Africa, 2018-2025



This chart shows difference in DTP3 coverage compared to 2019, by country.

Coverage was higher in 2025 compared to 2019 in 5 countries (CAR, Cameroon, Chad, Congo, and Equatorial Guinea).

The largest increase in coverage was 18% points in Chad from 53% in 2019 to 71% in 2025.

Coverage was lower than in 2019 in 4 countries (Burundi, DRC, Gabon, and Sao Tome and Principe).


The largest decline in coverage was 8% points in DRC from 73% in 2019 to 65% in 2025, Gabon from 70% in 2019 to 62% in 2025, and Sao Tome and Principe from 95% in 2019 to 87% in 2025.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: Dark grey line displays regional coverage difference for reference

ence compared to 2019, values above zero indicate coverage n percentage points higher than in 2019 and values below zero indicate coverage n percentage points lower than in 2019

DTP3 coverage, by country, Central Africa, 2019-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: The grey bar spans vaccine coverage across all years 2019-2025 and the dots represent coverage in specific years. Countries are arranged by high to low coverage in 2025 and coloured based on if coverage is lower (red), the same as (blue) or higher (green) than in 2019. An asterisk (*) indicates countries aiming to catch-up children missed since 2019 through a Big Catch-up Plan, supported by financing from The Gavi Alliance.


This chart shows the range of DTP3 coverage across years 2019 to 2025 (grey bars), and coverage in specific years (dots), by country. The chart can be used for assessing recovery to pre-pandemic levels and progress beyond 2019.

In 2024, 4 (out of 9) countries had lower coverage than in 2019.

In 2025, 0 countries had lower DTP3 coverage than in 2024.

In 2025, 4 countries had lower coverage than in 2019.

Between 2019 and 2025, DTP3 coverage increased in five countries and decreased in four countries.




This chart shows DTP3 coverage in 2019 and 2025, and change in coverage between the two years.

Overall, 4 out of 9 (44%) countries in Central Africa had a decline in coverage. The largest decline was in DRC, Gabon, and Sao Tome and Principe (-8 percentage points), followed by Burundi (-4 percentage points).

There were 5 (56%) countries with an increase in DTP3 coverage. Coverage remained the same in 0 countries between 2019 and 2025.

Between 2024 and 2025, DTP3 coverage increased in six countries, and no countries experienced a decline.




This chart shows DTP3 coverage in 2024 and 2025, and change in coverage between the two years.

Overall, 0 out of 9 (0%) countries in Central Africa had a decline in coverage.

There were 6 (67%) countries with an increase in DTP3 coverage. Coverage remained the same in 3 countries between 2024 and 2025.

In 2025, CAR and Gabon had the lowest DTP3 coverage, while DRC and Chad had the highest number of un- and undervaccinated children, highlighting the need to consider both coverage and absolute numbers to identify all vulnerable populations.



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Bars are ranked by ascending coverage.
Numbers in bubbles display top 10 rank based on absolute number of un- and undervaccinated children. Number in parentheses shows number of un- and undervaccinated children.

This chart shows DTP3 coverage in countries in Central Africa from lowest to highest coverage, and the rank of the top 10 countries with the most un- and undervaccinated children, based on absolute numbers.

In 2025, DTP3 coverage ranged from 46% in CAR to 89% in Burundi.


0 out of 9 (0%) countries had coverage of at least 90%.

DRC had DTP3 coverage of 65% and the highest absolute number of un- and undervaccinated children in the region, followed by Chad.

Large cohort countries may have high numbers of un- and undervaccinated children despite high vaccine coverage. It is advantageous to look at both coverage and absolute numbers of unvaccinated children to ensure vulnerable countries with small birth cohorts are not missed.

In 2025, 1 countries accounted for over half of all un- and under-vaccinated children in Central Africa, with DRC contributing the highest share.

DTP3 unvaccinated (% of total unvaccinated)



In 2025, 1 countries accounted for over half (70%) of all un- and under-vaccinated children in Central Africa.


DRC had the highest number of children who did not receive DTP3 (n=1.5 million), which is approximately 70% of all un- and under-vaccinated children in Central Africa (n=2.2 million).

The number of unvaccinated children is influenced by both the birth cohort size and programme performance. As a result, countries with large birth cohorts may still have a high number of unvaccinated children, even if coverage rates are high. At the same time, countries with high coverage will also have a greater absolute number of children who are vaccinated. Please refer to the heatmaps for country-specific coverage.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Number in legend corresponds to region ranking based on highest to lowest absolute number of unvaccinated children and reflects order of pie slices.

MCV1

MCV1 coverage remained relatively constant 61% in 2025, leaving 2.8 million children without any protection against measles. MCV2 coverage increased to 53% in 2025.



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: There may be discrepancies due to rounding.


Measles, because of its high transmissibility, acts as a 'canary in the coalmine', quickly exposing any immunity gaps in the population. The coverage of measles containing vaccine (MCV) is thus often used as a tracer for protection.

The percentage of children receiving MCV1 – typically at 9 or 12 months – remained relatively constant at 62%. This is similar to in 2019, where coverage was 63%.

2.8 million children missed their routine first dose of measles vaccine.

MCV2 is typically administered to children between 18 months and five years old. MCV2 coverage increased at 53% in 2025.

In 2025, MCV1 coverage was 62%: 4.4 million children received the measles vaccine, but 2.8 million remain unvaccinated against the highly contagious disease.



Sources: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision; United Nations, Department of Economic and Social Affairs, Population Division (2024). World Population Prospects 2024, Online Edition.

MCV1 coverage in 2025 (62%) was within 1% of coverage in 2019 (63%).

The number of children vaccinated with MCV1 increased 12% from 4 million in 2019 to 4.4 million in 2025.


The number of surviving infants increased approximately 14% from 6.3 million in 2019 to 7.2 million in 2025.

In 2025, 488,000 more children were vaccinated than in 2019.

In 2025, there were 907,000 more surviving infants (target population) than in 2019.

For vaccine coverage to increase, the number of children vaccinated must increase at a faster rate than the population increases.

MCV1 coverage, by country, Central Africa, 2000-2025




This heatmap shows MCV1 coverage, by region and year. Red indicates low coverage (<60%) and green indicates high coverage (≥90%). The changing colour gradient can be used to tracked changes in coverage over time and recovery from pandemic-related disruptions where applicable.


Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Countries ordered based on descending coverage in 2025

MCV1 coverage has improved since 2000 in all regions and exceeded pre-pandemic levels in most regions, but progress is uneven and gaps in unvaccinated children persist.

MCV1 coverage, Central Africa, 2000-2025




MCV1 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of infants unprotected against measles, Central Africa



In 2025, Central Africa ranked number 5 out of 5 AU regions, with MCV1 coverage of 62%.


MCV1 coverage (62%) was 1 percentage point lower than in 2019 (63%).

This equates to 2.8 million unvaccinated children in 2025 compared to 2.3 million unvaccinated children in 2019.


Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019.
There may be discrepancies in total Unvaccinated children (millions) due to rounding.

MCV1 coverage improved between 2024 and 2025 in most regions and exceeded pre-pandemic levels in most regions, but progress is uneven and gaps in unvaccinated children persist.

MCV1 coverage, Central Africa, 2018-2025




MCV1 coverage difference compared to 2019



Central Africa Eastern Africa Northern Africa Southern Africa Western Africa

Number of infants unprotected against measles, Central Africa




In 2025, Central Africa ranked number 5 out of 5 AU regions, with MCV1 coverage of 62%.

MCV1 coverage (62%) was 1 percentage point lower than in 2019 (63%).

This equates to 2.8 million unvaccinated children in 2025 compared to 2.3 million unvaccinated children in 2019.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Coverage difference compared to 2019 - values above zero indicate coverage higher than in 2019 and values below zero indicate coverage lower than in 2019.
There may be discrepancies in total Unvaccinated children (millions) due to rounding.

Trends in the number of measles cases and MCV coverage, 2012-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision;
 Reported measles and rubella cases and incidence rates by WHO Member States, as of 12 June 2026.
 Provisional data based on monthly data reported to WHO (Geneva) as of 12 June 2026.


This chart shows trends in the number of measles cases and MCV coverage (Central Africa).

In 2025, there were 10,981 confirmed measles cases - an increase from 10,575 in 2024.

Measles cases were reported in 8 countries (Burundi, Cameroon, CAR, Chad, Congo, DRC, Equatorial Guinea, and Gabon). DRC reported the highest number of cases (n=4,320), accounting for 39% of total cases in the region.

Very high vaccination coverage is necessary to interrupt measles transmission due to the virus's high transmissibility. The WHO recommends a minimum of 95% coverage with two doses of MCV through routine immunization services.

In 2025, 0 countries achieved at least 90% MCV1 coverage, while 5 had coverage below 70%.



MCV1 coverage (%) <60 60-69 70-79 80-89


Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: This map is stylized and based on an approximate scale. This map does not reflect a position by UNICEF or WHO on the legal status of any country or territory or the delimitation of any frontiers.

In Central Africa, 5 out of 9 (56%) countries had MCV1 coverage below 70% in 2025. These are CAR, Chad, DRC, Equatorial Guinea, and Gabon.

These countries account for approximately 5.6 million (78%) of the total annual cohort of surviving infants in the region.

There were 0 (0%) countries that achieved MCV1 of 90% or higher, accounting for <100 (0%) of the total cohort.

Coverage of MCV1, by country, Central Africa, 2018-2025



In 2025, Sao Tome and Principe had the highest coverage (87%), while CAR had the lowest coverage (44%).


MCV1 coverage increased in 4 out of 9 countries between 2024 and 2025. The largest increase was 9% points in Equatorial Guinea from 53% to 62%.

1 countries saw a decrease in coverage (Gabon). The largest decline was 1% points in Gabon from 57% to 56%. Coverage remained constant between 2024 and 2025 in the remaining 4 countries.

In 2025, 0 out of 9 countries had achieved MCV1 coverage of at least 90%.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Dark grey line displays regional coverage for reference

MCV1 coverage difference compared to 2019, by country, Central Africa, 2018-2025



This chart shows difference in MCV1 coverage compared to 2019, by country.

Coverage was higher in 2025 compared to 2019 in 5 countries (CAR, Cameroon, Chad, Congo, and Equatorial Guinea).

The largest increase in coverage was 28% points in Chad from 40% in 2019 to 68% in 2025.


Coverage was lower than in 2019 in 4 countries (Burundi, DRC, Gabon, and Sao Tome and Principe).

The largest decline in coverage was 10% points in DRC from 65% in 2019 to 55% in 2025.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: Dark grey line displays regional coverage difference for reference compared to 2019, values above zero indicate coverage n percentage points higher than in 2019 and values below zero indicate coverage n percentage points lower than in 2019

MCV1 coverage, by country, Central Africa, 2019-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Note: The grey bar spans vaccine coverage across all years 2019-2025 and the dots represent coverage in specific years. Countries are arranged by high to low coverage in 2025 and coloured based on if coverage is lower (red), the same as (blue) or higher (green) than in 2019. An asterisk (*) indicates countries aiming to catch-up children missed since 2019 through a Big Catch-up Plan, supported by financing from The Gavi Alliance.


This chart shows the range of MCV1 coverage across years 2019 to 2025 (grey bars), and coverage in specific years (dots), by country. The chart can be used for assessing recovery to pre-pandemic levels and progress beyond 2019.

In 2024, 5 (out of 9) countries had lower coverage than in 2019.

In 2025, 1 countries had lower MCV1 coverage than in 2024.

In 2025, 4 countries had lower coverage than in 2019.

In 2025, CAR and DRC had the lowest MCV1 coverage and were also among the countries with the highest number of unvaccinated children.



This chart shows MCV1 coverage in countries in Central Africa from lowest to highest coverage, and the rank of the top 10 countries with the most unvaccinated children, based on absolute numbers.

In 2025, MCV1 coverage ranged from 44% in CAR to 87% in Sao Tome and Principe.

0 out of 9 (0%) countries had coverage of at least 90%.


DRC had MCV1 coverage of 55% and the highest absolute number of unvaccinated children in the region, followed by Chad.

Large cohort countries may have high numbers of unvaccinated children despite high vaccine coverage. It is advantageous to look at both coverage and absolute numbers of unvaccinated children to ensure vulnerable countries with small birth cohorts are not missed.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Bars are ranked by ascending coverage.
Numbers in bubbles display top 10 rank based on absolute number of unvaccinated children. Number in parentheses shows number of unvaccinated children.

In 2025, 1 countries accounted for over half of all children unprotected against measles in Central Africa, with DRC contributing the highest share.

MCV1 unvaccinated (% of total unvaccinated)



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Number in legend corresponds to region ranking based on highest to lowest absolute number of unvaccinated children and reflects order of pie slices.

In 2025, 1 countries accounted for over half (71.4%) of all children unprotected against measles in Central Africa.



DRC had the highest number of children who did not receive MCV1 (n=2 million), which is approximately 71.4% of all children unprotected against measles in Central Africa (n=2.8 million).

The number of unvaccinated children is influenced by both the birth cohort size and programme performance. As a result, countries with large birth cohorts may still have a high number of unvaccinated children, even if coverage rates are high. At the same time, countries with high coverage will also have a greater absolute number of children who are vaccinated. Please refer to the heatmaps for country-specific coverage.

Childhood immunization: Additional charts

In 2025, DTP1 and DTP3/MCV1 drop-out was high in Central Africa

Vaccine drop-out, Central Africa, 2000-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Drop-out classification: <5% = low, 5-10% = medium, >10% = high

Drop-out rates show the % of children who received DTP1, but not DTP3/MCV1. Low drop-out rates indicate high retention of children in immunization programmes.


This chart shows trends in drop-out rates between DTP1 and DTP3, and DTP1 and MCV1.

In 2025, 16% of children who received DTP1 did not receive DTP3 (left), and 26% of children who received DTP1 did not receive MCV1 (right).

The high DTP drop-out rates imply poor ability to provide a complete series of vaccines early in life. The high DTP-MCV drop-out rates imply poor retention in immunization programmes and ability to provide a full course of vaccines in infancy (up to one year).

In 2025, Central Africa DTP drop-out was NA and DTP-MCV drop-out was NA than global drop-out rates, respectively.

Drop-out between DTP1 and DTP3 (%), by country, Central Africa, 2018-2025



This chart shows trends in drop-out rates between DTP1 and DTP3 by country.


In 2025, CAR had the highest drop-out with 22% of children who received DTP1 not receiving DTP3.

Burundi and Sao Tome and Principe had the lowest DTP drop-out (0%). Zero drop-out indicates that near enough all children that received DTP1 also received DTP3.

3 countries had DTP drop-out of more than 10%.

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Dark grey line displays regional dropout for reference
Drop-out classification: <5% = low, 5-10% = medium, >10% = high

Drop-out between DTP1 and MCV1 (%), by country, Central Africa, 2018-2025



This chart shows trends in drop-out rates between DTP1 and MCV1 by country.

In 2025, DRC had the highest drop-out with 33% of children who received DTP1 not receiving MCV1.


Sao Tome and Principe had the lowest DTP-MCV drop-out (0%).

6 countries had DTP-MCV drop-out of more than 10%.#n

There were 0 countries with negative DTP-MCV dropout (ie. more children received MCV1 than DTP1).

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Dark grey line displays regional dropout for reference
Drop-out classification: <5% = low, 5-10% = medium, >10% = high

Coverage of selected routine vaccines, Central Africa, 2000-2025




This chart shows coverage trends for 9 core routine vaccines recommended in childhood.

The steep increasing trends in coverage of IPV, MCV2, PCVC and RotaC is in-part related to increasing introduction in countries, along with improved coverage.

In 2025, MCV2 had the lowest coverage of all vaccines (53%), followed by IPVC (54%).

Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Breadth of Protection, Central Africa, 2000-2025



Breadth of protection (BOP) is a combination of the number of vaccines in a country's national immunization programme and the coverage achieved for each vaccine.

The radar chart displays the BOP (shaded area) and vaccine coverage (points), by year.


BOP increased from 12% in 2000 to 57% in 2025. This increase since 2000 is a result of incremental improvement in introducing new vaccines and expanding vaccination services to previously missed populations.

Enhancing BOP now depends partly on improving the coverage of HPV, RCV1, MCV2, IPVc, ROTAC, MCV1, PCVC, DTP3, HEPB3, HIB3, and IPV1 - vaccines with coverage less than 80% in 2025.

Breadth of protection = ((DTP3 x 3) + HepB3 + Hib3 + IPV1 + IPVc + MCV1 + MCV2 + PCV3 + RCV1 + RotaC + HPV) / 13

Breadth of protection with all recommended antigens has more than doubled since 2000, but reaching the next gains depends on closing coverage gaps in newer vaccines still below 57%.

Breadth of protection, Central Africa, 2000-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision

Breadth of protection (BoP) = ((DTP3 x 3) + HepB3 + Hib3 + IPV1 +IPVC + MCV1 + MCV2 + PCVC + RCV1 + RotaC + HPVc) / 13

Breadth of protection (BOP) is a combination of the number of vaccines in a country's national immunization programme and the coverage achieved for each vaccine.


The radar chart displays the BOP (shaded area) and vaccine coverage (points), by year.

BOP increased from 12% in 2000 to 57% in 2025. This increase since 2000 is a result of incremental improvement in introducing new vaccines and expanding vaccination services to previously missed populations.

Enhancing BOP now depends partly on improving the coverage of HPV, RCV1, MCV2, IPVc, ROTAC, MCV1, PCVC, DTP3, HEPB3, HIB3, and IPV1 - vaccines with coverage less than 80% in 2025.


Regional average coverage masks variation between countries across all vaccines, highlighting heterogeneity in immunization performance and the need for targeted, country-specific strategies to reach underserved populations.

Vaccine coverage (%), by country, Central Africa, 2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Countries are ranked within each region. Country ISO codes displayed. Regional average coverage is shown in the headers and by the dashed line.

Trends in number of countries with WUENIC estimates and coverage, Central Africa, 2011-2025



This chart coverage progress of selected vaccines alongside number of countries with introductions to aid understanding of progress.

WUENIC starts producing estimates from the first year a country reports data. For new vaccines, this is often the year of, or the year year after a vaccine in introduced. As such, the numbers serve as a proxy for year of introduction.

As WUENIC has evolved and SAGE recommendations updated, more vaccines have been added to the estimation process. Where estimates were started after 2011, the first year of estimation is indicated by the vertical dashed line.


The total number of countries that are recommended for each vaccine is indicated by the dashed horizontal line and corresponding value.

Countries without the vaccine are included in aggregates as 0.

HPV vaccination

Methods: • [Bruni et al. 2021, HPV vaccination introduction worldwide and WHO and UNICEF estimates of national HPV immunization coverage 2010–2019 \(supplementary materials\).](#)

In 2025, HPV last dose coverage among girls declined to 5% and coverage among boys remained the same at 4%



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Number of countries with HPV vaccination are indicated in parentheses. Countries without introduction are included in aggregates using a value of zero for the estimate.

HPV programme coverage is assessed according to the national schedule, and is used for short-term performance monitoring and SDG reporting.

In 2025, first dose (HPV1) coverage among girls declined to 5% and last dose (HPVc) coverage declined to 5%.


HPV1 coverage among boys remained the same at 4% and HPVc coverage remained the same at 4%.

HPV vaccine was available in 3* (out of 9) countries for girls and 1 for boys. Countries that have not introduced are included in aggregates with coverage of 0%.

Increasing coverage relies on reaching more girls and boys in existing programmes and introducing HPV vaccine in countries that have not yet introduced it.

*Includes partial introductions

HPVc programme coverage, girls, by country, Central Africa, 2010-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
Note: Only countries with HPV introduction and where estimates were made are displayed.
Countries without HPV vaccination are included in the aggregate with a value of 0.
Countries are ordered based on descending coverage in 2025.

The 'HPVc programme coverage' indicator is defined as the target population who received the last dose of HPV vaccine in the reporting year.


This heatmap shows HPVc programme coverage among girls, by country and year. The colour gradient is based on the best performing country.

In 2025, Sao Tome and Principe had the highest HPVc coverage (97%), and Central Africa had the lowest coverage (5%).

Countries without HPV vaccine in 2025 (n=7): Burundi, CAR, Chad, Congo, DRC, Equatorial Guinea, and Gabon.

SDG 3.b.1

SDG 3.b.1: Proportion of the target population covered by all vaccines included in their national programme, 2000-2025



Source: WHO/UNICEF Estimates of National Immunization Coverage, 2025 revision
 Note: The four vaccination coverage indicators contribute to SDG indicator 3.b.1 are: DTP3, MCV2, PCVC and HPVc
 The global target is 90% coverage of all four antigens by 2030

Four vaccination coverage indicators contribute to Sustainable Development Goal 3, indicator b.1: DTP3, PCVC, MCV2 and HPVc (females).

The IA2030 global target is 90% coverage of all four antigens by 2030.


In 2025, Central Africa had achieved at least 90% coverage of 0 out of the 4 vaccines.

Additional resources

[illegible]

	A	B	C	D	E	F	G	H	I	J
1	unicef_regio	isoc3	country	vaccine	year	target_grp	coverage	vaccinated	unvaccinated	target
2	ROSA	AFG	Afghanistan bcg		1982	births	10	57,000	514,000	571,000
3	ROSA	AFG	Afghanistan bcg		1983	births	10	56,000	503,000	539,000
4	ROSA	AFG	Afghanistan bcg		1984	births	11	62,000	501,000	563,000
5	ROSA	AFG	Afghanistan bcg		1985	births	17	100,000	490,000	591,000
6	ROSA	AFG	Afghanistan bcg		1986	births	18	107,000	489,000	596,000
7	ROSA	AFG	Afghanistan bcg		1987	births	27	161,000	434,000	595,000
8	ROSA	AFG	Afghanistan bcg		1988	births	40	238,000	358,000	596,000
9	ROSA	AFG	Afghanistan bcg		1989	births	38	233,000	380,000	614,000
10	ROSA	AFG	Afghanistan bcg		1990	births	30	191,000	446,000	637,000
11	ROSA	AFG	Afghanistan bcg		1991	births	21	134,000	505,000	640,000
12	ROSA	AFG	Afghanistan bcg		1992	births	19	127,000	541,000	668,000
13	ROSA	AFG	Afghanistan bcg		1993	births	17			760,000
14	ROSA	AFG	Afghanistan bcg		1994	births	15			852,000
15	ROSA	AFG	Afghanistan bcg		1995	births				899,000
16	ROSA	AFG	Afghanistan bcg		1996	births				932,000
17	ROSA	AFG	Afghanistan bcg		1997	births				962,000
18	ROSA	AFG	Afghanistan bcg		1998	births				986,000
19	ROSA	AFG	Afghanistan bcg		1999	births				1,013,000
20	ROSA	AFG	Afghanistan bcg		2000	births	30			1,037,000
21	ROSA	AFG	Afghanistan bcg		2001	births	43	455,000	596,000	1,007,000
22	ROSA	AFG	Afghanistan bcg		2002	births	46	467,000	548,000	1,015,000


UNICEF, WHO, World Bank, Gavi and
African Union regions



Additional immunization data resources can be found at:

<https://data.unicef.org/resources/immunization/>

*UNICEF, WHO, World Bank, Gavi and
African Union regions*



The collage features several distinct data visualizations:

- Top Left:** A line chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Top Right:** A line chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Middle Left:** A line chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Middle Right:** A line chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Bottom Left:** A bar chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Bottom Center:** A bar chart titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.
- Bottom Right:** A heatmap titled "Adverse Climate Impacts by Sector and Year, 2010-2025" showing the number of children vaccinated by sector (Health, Education, Agriculture, etc.) over time. The y-axis is labeled "Number of Children Vaccinated" and ranges from 0 to 100. The x-axis is labeled "Year" and ranges from 2010 to 2025.

[illegible]

Interactive
WUENIC
country profiles
can be found at:

<https://worldhealthorg.shinyapps.io/wuenic-trends/>

Short feedback questionnaire

(5 minutes)

We are seeking your feedback on the global groupings (GAVI, African Union, World Bank Income, WHO and UNICEF) and country-level PowerPoint slides developed for the release of global immunization estimates. Your input will help us understand their usefulness and identify areas for improvement.

Please take a few moments to complete this short survey and have your voice heard:

<https://forms.office.com/e/Qv1HXxxNZQ>

