

WUENIC Talking Points: Comoros



Revision Year: 2025

Contents

Overview	2
DTP1: Reaching zero-dose children	3
DTP3: A marker of routine immunization service delivery to children	4
MCV: Canary in the coal mine	5
HPV	6
Regional Comparison	7
Additional Vaccines	8
Appendix	9


Comoros in the 'Africa' region



Overview


- None of the 10 childhood vaccines tracked for Comoros achieved coverage of 90% or higher in 2025.
- Coverage of the first dose of diphtheria-tetanus-pertussis-containing vaccine (DTP1) was the same in 2025 as it was in 2024 (80%), coverage of the third dose of DTP-containing vaccine (DTP3) was the same in 2025 as it was in 2024 (75%), and coverage of the first dose of the measles-containing vaccine (MCV1) was the same in 2025 as it was in 2024 (70%) in 2025.
- In 2025, the number of zero-dose children for DTP in Comoros (5,000) was approximately 49% higher than the annual number proposed to achieve the IA2030 target of halving zero-dose children for DTP by 2030 (3,000), based on a linear trajectory of decline.

Vaccine coverage (%), Comoros, 2000–2025



Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

Estimated number of zero-dose children, 2000–2025 and target by 2030, Comoros




Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.
 Note: The Immunization Agenda 2030 (IA2030) calls on all countries to reduce the number of zero-dose children in 2019 in half by 2030.
 Dark blue bars are the estimated number of zero-dose children in 2000–2025, light blue bar is the target number of zero-dose children by 2030.
 The line and points show the yearly progress and trajectory to meet the target by 2030, based on a linear decline.

DTP1: Reaching zero-dose children

- Coverage of DTP1 stood at 80% in 2025. This is lower than in 2019 (81%) and the same as in 2024 (80%).
- The percentage of zero-dose children was 49 percentage points higher than the IA2030 target in 2025. There were approximately 5,000 zero-dose children in 2025, more than the 4,000 in 2019 and more than the 5,000 in 2024.
- Coverage of DTP1 in Comoros (80%) was lower than the Eastern and Southern Africa regional average of 85% in 2025.
- DTP1-to-DTP3 dropout in 2025 was 6%, 2 percentage points higher than in 2019 (4%). Medium DTP dropout rates imply moderate ability to provide a complete vaccine series early in life.

DTP1 coverage and zero-dose children over time, Comoros, 2000–2025




Dotted line shows the IA2030 target of 90% coverage
Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

DTP3: A marker of routine immunization service delivery to children

- Coverage of DTP3 stood at 75% in 2025. This is lower than in 2019 (78%) and the same as in 2024 (75%).
- The number of children vaccinated with DTP3 decreased from 18,000 in 2019 to 18,000 in 2025.
- In 2025, there were 6,000 children un- or under-vaccinated in Comoros.
- In 2025, 6% of children who received DTP1 did not receive DTP3. This 6% dropout rate was 2 percentage points higher than in 2019 (4%) and the same as the 2024 dropout rate (6%).
- Coverage of DTP3 in Comoros (75%) was lower than the Eastern and Southern Africa regional average of 80% in 2025.


DTP3 coverage and un- or under-vaccinated children over time, Comoros, 2000–2025



Dotted line shows the IA2030 target of 90% coverage
Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

MCV: Canary in the coal mine


- Coverage of MCV1 stood at 70% in 2025. This is lower than in 2019 (76%) and the same as in 2024 (70%).
- Coverage of the second dose of the measles-containing vaccine (MCV2) NA. This leaves 7,000 children without any protection against measles and another <500 with only partial protection.
- In 2025, 12% of children who received DTP1 did not receive MCV1. This 12% dropout rate was 6 percentage points higher than in 2019 (6%) but 1 percentage points lower than the 2024 dropout rate (13%).
- Comoros has had MCV1 coverage below 80% for all of the last 3 years.
- Coverage of MCV1 in Comoros (70%) was lower than the Eastern and Southern Africa regional average of 77% in 2025.



Dotted line shows the IA2030 target of 90% coverage
 Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

HPV

- In 2025, programme coverage among females was 26% for the first dose (HPV1) and 26% for the final dose (HPVc).
- NA
- In 2025, approximately 7,000 females in Comoros did not receive any HPV vaccination (un-vaccinated).
- Among Eastern and Southern Africa region peers, Comoros ranked 16 out of 17 countries in HPV1 coverage in 2025.




Dotted line shows the IA2030 target of 90% coverage
Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

Regional Comparison

The chart below shows how Comoros compares to other countries in the Eastern and Southern Africa region on DTP1, DTP3, MCV1, and HPVc coverage in 2025. The dashed line represents the regional average for each vaccine.

Coverage comparison: Comoros vs. ESAR peers, 2025

Dashed line = regional average




Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

Additional Vaccines

- Coverage of BCG stood at 84% in 2025. This is lower than in 2019 (87%) and lower than in 2024 (85%).
- Coverage of HepB3 stood at 75% in 2025. This is lower than in 2019 (78%) and the same as in 2024 (75%).
- Coverage of Hib3 stood at 75% in 2025. This is lower than in 2019 (78%) and the same as in 2024 (75%).
- Coverage of IPV1 stood at 75% in 2025. This is higher than in 2019 (72%) and the same as in 2024 (75%).

Additional vaccine coverage (%), Comoros, 2000–2025



Source: WHO/UNICEF estimates of national immunization coverage, 2025 revision.

Appendix

WHO/UNICEF Estimates of National Immunization Coverage (WUENIC)

Each year, WHO and UNICEF review national immunization data from Member States, including administrative and official coverage, survey estimates and contextual information, and use rules-based data triangulation approach to assess coverage. Estimates are produced independently, without borrowing data from other countries, and are not based on ad hoc adjustments. In some cases, they rely on a single source, typically nationally reported coverage. When data for a specific country-vaccine-year are unavailable, values from adjacent years are interpolated or extrapolated, and conflicting sources are assessed to determine the most reliable estimate, considering potential biases and expert opinion.

Methods:

- [Burton et al. 2009. WHO and UNICEF estimates of national infant immunization coverage: methods and processes](#)
- [Burton et al. 2012. A formal representation of the WHO and UNICEF estimates of national immunization coverage: a computational logic approach](#)
- [Brown et al. 2013. An introduction to the grade of confidence used to characterize uncertainty around the WHO and UNICEF estimates of national immunization coverage](#)

These talking points present the latest WUENIC estimates (published 15 July 2026).

Definitions of immunization terms:

- Bacillus Calmette-Guerin (BCG): vaccine against tuberculosis
- Hepatitis B birth dose, given within 24 hours after birth (HepBB)
- Diphtheria, tetanus, and pertussis vaccine, first dose (DTP1) and third dose (DTP3)
- Hepatitis B vaccine, third dose (HepB3)
- Haemophilus influenzae type B vaccine, third dose (Hib3)
- Inactivated polio vaccine, first dose (IPV1) and last dose (IPVC)
- Measles containing vaccine, first dose (MCV1) and second dose (MCV2)
- Rotavirus vaccine, last dose (RotaC)
- Pneumococcal vaccine, last dose (PCV)
- Yellow Fever vaccine (YFV)
- Meningococcal A vaccine (MengA)
- Human Papillomavirus vaccine, first dose (HPV1) and last dose (HPVc)

Additional resources: <https://data.unicef.org/resources/immunization/>