




DEMOGRAPHICS

Total population (000)	10,943	(2011)
Total under-five population (000)	1,912	(2011)
Births (000)	449	(2011)
Birth registration (%)	63	(2010)
Total under-five deaths (000)	23	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	21	(2011)
Infant mortality rate (per 1000 live births)	38	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,500	(2010)
Lifetime risk of maternal death (1 in N)	54	(2010)
Total fertility rate (per woman)	5.3	(2011)
Adolescent birth rate (per 1000 women)	41	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	71	(2010)
Low birthweight incidence (moderate and severe, %)	7	(2010)	Introduction of solid, semi-solid/soft foods (%)	79	(2010)
			Vitamin A two dose coverage (%)	76	(2011)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

