

Nutrition Profile United Arab Emirates

DEMOGRAPHICS


Total population (000)	4,485	(2008)
Total under-five population (000)	307	(2008)
Total number of births (000)	63	(2008)
Under-five mortality rate (per 1,000 live births)	8	(2008)
Total number of under-five deaths (000)	0	(2008)
Infant mortality rate (per 1,000 live births)	7	(2008)
Neonatal mortality rate (per 1,000 live births)	4	(2004)
HIV prevalence rate (15-49 years, %)		-
Population below international poverty line of US\$1.25 per day (%)		





Other NS

NUTRITIONAL STATUS



Other NS

INFANT AND YOUNG CHILD FEEDING

100 %

80

40 60

0 20

Source: Other NS 1995



United Arab Emirates







ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE

Pregnancy	Birth	0-5 months	6-23 months		24-59 months
Use of iron-folic acid	f Breastmilk Substitutes	Partial			
supplements	(within 1 hour of birth)	Maternity protection in accordance with ILO Convention 183			
Household consumption of . adequately iodized salt	Infants not weighed at birth ·	Exclusive breastfeeding 34% (<6 months)	Timely introduction of complementary foods (with continued breastfeeding)	52%	
		Continued breastfeeding at two years	29%		
To increase childre	Full coverage of vitamin A supplementation				
development and p need to be delivere	National guidelines for management of severe acute malnutrition incorporating the community-based approach				
first two years of t	ne chila s lite.	Policy on new ORS formula and zinc for management of diarrhoea*			
	Informa	Policy on community treatment of pneumonia with antibiotics			

MATERNAL NUTRITION/HEALTH

WATER AND SANITATION





Diarrhoea:

Shared facility

Open defecation

98

Urhan

4%

95

Rural

Source: WHO/UNICEF JMP, 2010

10% Pneumonia:

DISPARITIES IN NUTRITION

		Gender			Residence			Wealth quintile					
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)					-	-					-		
Underweight prevalence (WHO Child Growth Standards, %)											-		
Wasting prevalence (WHO Child Growth Standards, %)		-	-		-	-					-	-	
Infants not weighed at birth (%)		-	-				-			-	-		
Early initiation of breastfeeding (%)		-	-	-	-	-	-			-	-	-	•
Women with low BMI ($<$ 18.5 kg/m², %)		-			-	-					-	-	