

Nutrition Profile

Bulgaria

DEMOGRAPHICS

Total population (000)	7,593	(2008)
Total under-five population (000)	349	(2008)
Total number of births (000)	73	(2008)
Under-five mortality rate (per 1,000 live births)	11	(2008)
Total number of under-five deatl	1	(2008)
Infant mortality rate (per 1,000	9	(2008)
Neonatal mortality rate (per 1,000 live	7	(2004)
HIV prevalence rate (15-49 years,	-	-
Population below international poverty line of US\$1.25 per day	<2	(2003)





NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards	Stunted (under-fives, 000):	- Underweight (under-fives, 000):	-
	Share of developing world stunting burden	- Wasted (under-fives, 000):	
	Stunting country rank:	- Severely wasted (under-fives, 000):	-
Current nutritional status Percentage of children < 5 years old suffering from: No Data	Stunting trends Percentage of children < 5 years old stunted NCHS reference population No Data	Underweight trends Percentage of children < 5 years old underweight, NCHS reference population No Data	

INFANT AND YOUNG CHILD FEEDING

Infant feeding practices, by age	
	Weaned (not Breastfed and non-milk liquids Breastfed and solid/ semi-solid foods Breastfed and plain water only Breastfed and other milk/formula Exclusively breastfed
No Data	Exclusive breastfeeding Percentage of infants < 6 months old exclusively breastfed
	No Data

Bulgaria

MICRONUTRIENTS





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH

WATER AND SANITATION





Diarrhoea:



DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	icator Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Underweight prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Wasting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Infants not weighed at birth (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Early initiation of breastfeeding (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Women with low BMI (<18.5 kg/m ² , %)	-	-	-	-	-	-	-	-	-	-	-		-