

Belarus

DEMOGRAPHICS

Total population (000)	9,679	(2008)
Total under-five population (000)	472	(2008)
Total number of births (000)	96	(2008)
Under-five mortality rate (per 1,000 live births)	13	(2008)
Total number of under-five deatl	1	(2008)
Infant mortality rate (per 1,000	11	(2008)
Neonatal mortality rate (per 1,000 live	3	(2004)
HIV prevalence rate (15-49 years,	0.2	(2007)
Population below international poverty line of US\$1.25 per day	<2	(2005)






NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards


 Stunted (under-fives, 000):
 21
 Underweight (under-fives, 000):
 6

 Share of developing world stunting burden
 0.0
 Wasted (under-fives, 000):
 10


 Stunting country rank:
 115
 Severely wasted (under-fives, 000):
 3








INFANT AND YOUNG CHILD FEEDING






Belarus

MICRONUTRIENTS

Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year


No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION







Under-five deaths caused Diarrhoea:

0% Pneumonia:

DISPARITIES IN NUTRITION

Indicator	Gender			Residence			Wealth quintile						
	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poores	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	5	4	1.3	3	8	0.4	10	5	3	3	2	0.2	MICS 2005
Underweight prevalence (WHO Child Growth Standards, %)	2	1	2.0	1	2	0.5	2	1	1	2	0	-	MICS 2005
Wasting prevalence (WHO Child Growth Standards, %)	3	2	1.5	2	2	1.0	2	2	1	3	2	1.0	MICS 2005
Infants not weighed at birth (%)	-	-	-	1	1	1.0	2	0	0	0	1	0.5	MICS 2005
Early initiation of breastfeeding (%)	-	-	-	23	18	1.3	17	17	15	22	30	1.8	MICS 2005
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-