



DEMOGRAPHICS

Total population (000)	687	(2008)
Total under-five population (000)	71	(2008)
Total number of births (000)	15	(2008)
Under-five mortality rate (per 1,000 live births)	81	(2008)
Total number of under-five deatl	1	(2008)
Infant mortality rate (per 1,000	54	(2008)
Neonatal mortality rate (per 1,000 live	30	(2004)
HIV prevalence rate (15-49 years,	0.1	(2007)
Population below international poverty line of US\$1.25 per day	26	(2003)






NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards


 Stunted (under-fives, 000):
 34
 Underweight (under-fives, 000):
 10

 Share of developing world stunting burden
 0.0
 Wasted (under-fives, 000):
 2

 Stunting country rank:
 110
 Severely wasted (under-fives, 000):
 1









INFANT AND YOUNG CHILD FEEDING




Bhutan

MICRONUTRIENTS








* Estimates may not be comparable.


ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births) 440	(2005)
Maternal mortality ratio, reported (per 100,000 live births) 260	(2000)
Total number of maternal deaths 280	(2005)
Lifetime risk of maternal deaths (1 in :) 55	(2005)
Women with low BMI (< 18.5 kg/m², -	-
Anaemia, non-pregnant woman (<120 g/l, 55	(2002)
Antenatal care (at least one visit, %) 88	(2007)
Antenatal care (at least four visits, %)	-
Skilled attendant at birth (%) 71	(2007)
Low birthweight (<2,500 grams, %) 15	(1999)
Primary school net enrolment or attendance ratio (% female, % male) 67, 74	(2003)
Gender parity index (primary school net 0.91 enrolment or attendance ratio)	(2003)





Under-five deaths caused

Diarrhoea:

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	52	43	1.2	-	-	-	-	-	-	-	-	-	Other NS 1999
Underweight prevalence (WHO Child Growth Standards, %)	17	12	1.4	-	-	-	-	-	-	-	-	-	Other NS 1999
Wasting prevalence (WHO Child Growth Standards, %)	3	2	1.5	-	-	-	-	-	-	-	-	-	Other NS 1999
Infants not weighed at birth (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Early initiation of breastfeeding (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-