

Dominica

DEMOGRAPHICS

Total population (000) 67 (2008)			
Total number of births (000) 1 (2008) Under-five mortality rate (per 1,000 live births) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) HIV prevalence rate (15-49 years, %) Population below international	Total population (000)	67	(2008)
Under-five mortality rate (per 1,000 live births) Total number of under-five deaths (000) Infant mortality rate (per 1,000 live births) Neonatal mortality rate (per 1,000 live births) HIV prevalence rate (15-49 years, %) Population below international	Total under-five population (000)	3	(2008)
births) 11 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 10 (2004) HIV prevalence rate (15-49 years, %)	Total number of births (000)	1	(2008)
Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 10 (2004) HIV prevalence rate (15-49 years, %)		11	(2008)
Neonatal mortality rate (per 1,000 live births) 10 (2004) HIV prevalence rate (15-49 years, %)	Total number of under-five deaths (000)	0	(2008)
births) 10 (2004) HIV prevalence rate (15-49 years, %) Population below international	Infant mortality rate (per 1,000 live births)	9	(2008)
Population below international	• •	10	(2004)
•	HIV prevalence rate (15-49 years, %)		-
	•		





NUTRITIONAL STATUS

Burden of undernutrition (2008)
WHO Child Growth Standards
Stunted (under-fives, 000):
Share of developing world stunting burden (%):
Stunting country rank:

Stunted (under-fives, 000):
Stunting country rank:

Underweight (under-fives, 000):
Wasted (under-fives, 000):
Severely wasted (under-fives, 000):

Current nutritional status

Percentage of children < 5 years old suffering

No Data

Stunting trends

Percentage of children < 5 years old stunted NCHS reference population

No Data

Underweight trends

Percentage of children < 5 years old underweight NCHS reference population

No Data

INFANT AND YOUNG CHILD FEEDING

Dominica


MICRONUTRIENTS

Vitamin A supplementation Percentage of children 6-59 months old receiving two of vitamin A during calendar year

No Data

lodized salt consumption trends Percentage of households consuming adequately iodized salt

No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE


Pregnancy	Birth	0-5 months	6-23 months	>	24-59 months		
Use of iron-folic acid	of iron-folic acid Early initiation of breastfeeding International Code of Marketing of Br						
supplements .	(within 1 hour of birth)	ce with ILO Convention 183	No				
Household consumption of adequately iodized salt	Infants not weighed at birth .	Exclusive breastfeeding (< 6 months)	· · · · · · · · · · · · · · · · · · ·				
		Continued breastfeeding at two years					
To increase children's chances of survival, improve development and prevent stunting, nutrition interventions need to be delivered during the mother's pregnancy and the			Full coverage of vitamin A supplementation				
			National guidelines for management of severe acute malnutrition incorporating the community-based approach				
first two years of the	Policy on new ORS formula and zinc for management of diarrhoea*						
	Policy on community treatment of pneumonia with antibiotics*						

MATERNAL NUTRITION/HEALTH

WATER AND SANITATION







Under-five deaths caused by:

Diarrhoea:

0% Pneumonia:

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-				-	-	-						
Underweight prevalence (WHO Child Growth Standards, %)	-				-		-						•
Wasting prevalence (WHO Child Growth Standards, %)			-		-	-	-						
Infants not weighed at birth (%)			-		-								
Early initiation of breastfeeding (%)			-		-	-	-						
Women with low BMI ($<$ 18.5 kg/m², %)	-	-	-				-		-				