

Dominican Republic

DEMOGRAPHICS

Total population (000)	9,953	(2008)
Total under-five population (000)	1,086	(2008)
Total number of births (000)	224	(2008)
Under-five mortality rate (per 1,000 live births)	33	(2008)
Total number of under-five deatl	7	(2008)
Infant mortality rate (per 1,000	27	(2008)
Neonatal mortality rate (per 1,000 live	18	(2004)
HIV prevalence rate (15-49 years,	1.1	(2007)
Population below international poverty line of US\$1.25 per day	5	(2005)








NUTRITIONAL STATUS

Burden of undernutrition (2008) WHO Child Growth Standards


Stunted (under-fives, 000):	197	Underweight (under-fives, 000):	77
Share of developing world stunting burden	0.1	Wasted (under-fives, 000):	29
Stunting country rank:	74	Severely wasted (under-fives, 000):	11








INFANT AND YOUNG CHILD FEEDING






Dominican Republic


MICRONUTRIENTS

Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data






ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births)	150	(2005)
Maternal mortality ratio, reported (per 100,000 live births)	160	(1997- 2007)
Total number of maternal deaths	310	(2005)
Lifetime risk of maternal deaths (1 in :)	230	(2005)
Women with low BMI (< 18.5 kg/m²,	-	-
Anaemia, non-pregnant woman (<120 g/	ı, 2 7	(2008)
Antenatal care (at least one visit, %)	99	(2007)
Antenatal care (at least four visits, %)	95	(2007)
Skilled attendant at birth (%)	98	(2007)
Low birthweight (<2,500 grams, %)	11	(2007)
attendance ratio (% female, % male)	, 88	(2007)
Gender parity index (primary school net enrolment or attendance ratio)	1.02	(2007)





Under-five deaths caused by:	Diarrhoea:	11%	10%

DISPARITIES IN NUTRITION

	Gender			Residence		Wealth quintile							
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poores	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	22	14	1.6	-	-	-	23	19	7	-	-	-	DHS 2007
Underweight prevalence (WHO Child Growth Standards, %)	10	5	2.0	-	-	-	11	5	1	-	-	-	DHS 2007
Wasting prevalence (WHO Child Growth Standards, %)	3	2	1.5	-	-	-	5	5	4	-	-	-	DHS 2007
Infants not weighed at birth (%)	-	-	-	3	5	0.6	7	3	3	2	2	0.3	DHS 2007
Early initiation of breastfeeding (%)	73	75	1.0	-	-	-	76	71	73	-	-	-	DHS 2007
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-