

Grenada

DEMOGRAPHICS

Total population (000)	104	(2008)
Total under-five population (000)	9	(2008)
Total number of births (000)	2	(2008)
Under-five mortality rate (per 1,000 live births)	15	(2008)
Total number of under-five deatl	0	(2008)
Infant mortality rate (per 1,000	13	(2008)
Neonatal mortality rate (per 1,000	11	(2004)
live HIV prevalence rate (15-49 years,	-	-
Population below international poverty line of US\$1.25 per day	•	-





NUTRITIONAL STATUS

Burden of undernutrition (2008) WHO Child Growth Standards

Stunted (under-fives, 000): Share of developing world stunting burden Stunting country rank:

Underweight (under-fives, 000): - Wasted (under-fives, 000):

- Severely wasted (under-fives, 000):

Current nutritional status

Percentage of children < 5 years old suffering from:

No Data

Stunting trends Percentage of children < 5 years old stunted NCHS reference population

No Data

Underweight trends

Percentage of children < 5 years old underweight, NCHS reference population


No Data

INFANT AND YOUNG CHILD FEEDING

Infant feeding practices, by age

No Data

Breastfed and non-milk liquids Weaned (not Breastfed and solid/ Breastfed and plain semi-solid foods water only Breastfed and other Exclusively breastfed milk/formula




Grenada

MICRONUTRIENTS

Vitamin A supplementation

Percentage of children 6-59 months old receiving t two doses of vitamin A during calendar year


No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION







Under-five deaths caused by: Diarrhoea: 1% 6%

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Underweight prevalence (WHO Child Growth Standards, %)	-	-	•	-	-	-	-	-	-	-	-	-	-
Wasting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Infants not weighed at birth (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Early initiation of breastfeeding (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-