


Guyana

DEMOGRAPHICS








NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards


 Stunted (under-fives, 000):
 12
 Underweight (under-fives, 000):
 7

 Share of developing world stunting burden
 0.0
 Wasted (under-fives, 000):
 6


 Stunting country rank:
 120
 Severely wasted (under-fives, 000):
 2








INFANT AND YOUNG CHILD FEEDING





Guyana

MICRONUTRIENTS


Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data

lodized salt consumption trends
Percentage of households consuming
adequately iodized salt
14,000 newborns are unprotected
against IDD (2008)

No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births) 47	70	(2005)
Maternal mortality ratio, reported (per 100,000 live births)	60	(2005)
Total number of maternal deaths 7	73	(2005)
Lifetime risk of maternal deaths (1 in :)	90	(2005)
Women with low BMI (< 18.5 kg/m²,	-	-
Anaemia, non-pregnant woman (<120 g/l, 5	54	(1996-
Antenatal care (at least one visit, %)	31	(2006)
Antenatal care (at least four visits, %)	-	-
Skilled attendant at birth (%)	33	(2006)
Low birthweight (<2,500 grams, %)	9	(2006-
Primary school net enrolment or attendance ratio (% female, % male)	96	(2006)
Gender parity index (primary school net enrolment or attendance ratio)	1	(2006)





Under-five deaths caused by:

Diarrhoea: 12

7%

DISPARITIES IN NUTRITION

Indicator	Gender			Residence			Wealth quintile						
	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poores	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	18	16	1.1	14	18	0.8	28	14	10	11	12	0.4	MICS 2006- 2007
Underweight prevalence (WHO Child Growth Standards, %)	10	9	1.1	10	9	1.1	10	12	9	9	4	0.4	MICS 2006- 2007
Wasting prevalence (WHO Child Growth Standards, %)	8	9	0.9	12	7	1.7	4	11	10	12	5	1.3	MICS 2006- 2007
Infants not weighed at birth (%)	-	-	-	27	21	1.3	30	23	14	21	17	0.6	MICS 2006-
Early initiation of breastfeeding (%)	-	-	-	42	44	1.0	45	43	47	42	34	0.8	MICS 2006-
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-