


Mexico

DEMOGRAPHICS








NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards


 Stunted (under-fives, 000):
 1,594
 Underweight (under-fives, 000):
 350

 Share of developing world stunting burden
 0.8
 Wasted (under-fives, 000):
 206


 Stunting country rank:
 22
 Severely wasted (under-fives, 000):








INFANT AND YOUNG CHILD FEEDING






Mexico

MICRONUTRIENTS

Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births) 60	(2005)
Maternal mortality ratio, reported (per 100,000 live births) 56	(2007)
Total number of maternal deaths 1,300	(2005)
Lifetime risk of maternal deaths (1 in :) 670	(2005)
Women with low BMI (< 18.5 kg/m², -	-
Anaemia, non-pregnant woman (<120 g/l, 16	(2006)
Antenatal care (at least one visit, %) 94	(2006)
Antenatal care (at least four visits, %) -	-
Skilled attendant at birth (%) 93	(2006)
Low birthweight (<2,500 grams, %) 8	(2006)
Primary school net enrolment or attendance ratio (% female, % male) 97, 98	(2006)
Gender parity index (primary school net 0.99 enrolment or attendance ratio)	(2006)





Under-five deaths caused by: Diarrhoea: 4% 9%

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Underweight prevalence (WHO Child Growth Standards, %)	-	-	•	-	-	-	-	-	-	-	-	-	
Wasting prevalence (WHO Child Growth Standards, %)	-	-	-	-	-	-	-	-	-	-	-	-	-
Infants not weighed at birth (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Early initiation of breastfeeding (%)	-	-	-	-	-	-	-	-	-	-	-	-	-
Women with low BMI (<18.5 kg/m², %)	-	-	-	-	-	-	-	-	-	-	-	-	-