


Korea, Democratic People's Republic of

Nutrition Profile

DEMOGRAPHICS










NUTRITIONAL STATUS

Burden of undernutrition (2008) WHO Child Growth Standards


Stunted (under-fives, 000):	704	Underweight (under-fives, 000):	280
Share of developing world stunting burden (%):	0.4	Wasted (under-fives, 000):	137
Stunting country rank:	43	Severely wasted (under-fives, 000):	55








INFANT AND YOUNG CHILD FEEDING



No Data





2004 Other NS


20

Korea, Democratic People's Republic of

MICRONUTRIENTS








* Estimates may not be comparable.


ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH

WATER AND SANITATION







Under-five deaths caused by:

Diarrhoea:

26% Pneumonia:

DISPARITIES IN NUTRITION

Indicator	Gender			Residence			Wealth quintile						
	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	47	42	1.1	45	44	1.0	-	-	-	-			Other NS 2002
Underweight prevalence (WHO Child Growth Standards, %)	19	16	1.2	17	19	0.9	-						Other NS 2002
Wasting prevalence (WHO Child Growth Standards, %)	9	8	1.1	9	9	1.0	-		-	-			Other NS 2002
Infants not weighed at birth (%)							-						
Early initiation of breastfeeding (%)	-	-		-			-	-	-	-	-	-	
Women with low BMI (< 18.5 kg/m², %)	-	-					-	-	-	-			