

Nutrition Profile

Sudan

DEMOGRAPHICS

41,348	(2008)
5,836	(2008)
1,296	(2008)
109	(2008)
138	(2008)
70	(2008)
27	(2004)
1.4	(2007)
	5,836 1,296 109 138 70 27





NUTRITIONAL STATUS

Burden of undernutrition (2008) WHO Child Growth Standards

Stunted (under-fives, 000):	2,305	Underweight (under-fives, 000):	1,576
Share of developing world stunting burden (%):	1.2	Wasted (under-fives, 000):	945
Stunting country rank:	15	Severely wasted (under-fives, 000):	403

Current nutritional status
Percentage of children < 5 years old suffering
from:

Stunting

40

33

Underweight

Wasting


16


WHO Child Growth Standards

NCHS reference population


0 20 40 60 80 100 %


Source: Other NS 2006






INFANT AND YOUNG CHILD FEEDING







Sudan

MICRONUTRIENTS







* Estimates may not be comparable


ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH

WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births)	450	(2005)
Maternal mortality ratio, reported (per 100,000 live births)	1,100	(2006)
Total number of maternal deaths	5,300	(2005)
Lifetime risk of maternal deaths (1 in :)	53	(2005)
Women with low BMI ($< 18.5 \mathrm{kg/m^2}$, %)		-
Anaemia, non-pregnant woman (< 120 g/l, %)	43	(2008)
Antenatal care (at least one visit, %)	64	(2006)
Antenatal care (at least four visits, %)		
Skilled attendant at birth (%)	49	(2006)
Low birthweight (< 2,500 grams, %)	31	(1999)
Primary school net enrolment or attendance ratio (% female, % male)	52, 56	(2006)
Gender parity index (primary school net enrolment or attendance ratio)	0.93	(2006)





15%

13% Pneumonia:

DISPARITIES IN NUTRITION

Indicator	Gender			Residence			Wealth quintile						
	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	42	37	1.1	35	42	0.8	39	45	44	39	28	0.7	Other NS 2006
Underweight prevalence (WHO Child Growth Standards, %)	28	26	1.1	21	30	0.7	31	33	30	23	17	0.5	Other NS 2006
Wasting prevalence (WHO Child Growth Standards, %)	17	15	1.1	14	18	0.8	24	20	15	12	11	0.5	Other NS 2006
Infants not weighed at birth (%)	-	-			-			-					
Early initiation of breastfeeding (%)	-	-	-		-		-	-					
Women with low BMI ($<$ 18.5 kg/m², %)	-	-			-							•	