

Solomon Islands

DEMOGRAPHICS

Total population (000)	511	(2008)
Total under-five population (000)	73	(2008)
Total number of births (000)	16	(2008)
Under-five mortality rate (per 1,000 live births)	36	(2008)
Total number of under-five deaths (000)	1	(2008)
Infant mortality rate (per 1,000 live births)	30	(2008)
Neonatal mortality rate (per 1,000 live births)	23	(2004)
HIV prevalence rate (15-49 years, %)		
Population below international poverty line of US\$1.25 per day (%)		







NUTRITIONAL STATUS


Burden of undernutrition (2008) WHO Child Growth Standards

Stunted (under-fives, 000):	24	Underweight (under-fives, 000):	9
Share of developing world stunting burden (%):	0.0	Wasted (under-fives, 000):	3
Stunting country rank:	113	Severely wasted (under-fives, 000):	1



Infant feeding practices, by age






INFANT AND YOUNG CHILD FEEDING

No Data





Solomon Islands

MICRONUTRIENTS


Vitamin A supplementation

Percentage of children 6-59 months old receiving

No Data

lodized salt consumption trends*
Percentage of households consuming adequately iodized salt

No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE



MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births)	220	(2005)
Maternal mortality ratio, reported (per 100,000 live births)	140	(1999)
Total number of maternal deaths	34	(2005)
Lifetime risk of maternal deaths (1 in :)	100	(2005)
Women with low BMI ($< 18.5 \ kg/m^2, \%)$	-	-
Anaemia, non-pregnant woman (< 120 g/l, %)	39	(2008)
Antenatal care (at least one visit, %)	74	(2007)
Antenatal care (at least four visits, %)	65	(2007)
Skilled attendant at birth (%)	70	(2007)
Low birthweight ($<$ 2,500 grams, %)	13	(2007)
Primary school net enrolment or attendance ratio (% female, % male)	62, 62	(2005)
Gender parity index (primary school net enrolment or attendance ratio)	1	(2005)





Under-five deaths caused by: Diarrhoea: 8% Pneumonia: 8%

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	36	32	1.1	-	-	÷	-	-	-		-	-	Other NS 1989
Underweight prevalence (WHO Child Growth Standards, %)	18	14	1.3	-		•	-		-		-		Other NS 1989
Wasting prevalence (WHO Child Growth Standards, %)	8	6	1.3	-	-	-	-	-	-		-	-	Other NS 1989
Infants not weighed at birth (%)				7	20	0.4	30	20	18	14	8	0.3	DHS 2007
Early initiation of breastfeeding (%)	-		-	78	75	1.0	72	79	76	75	73	1.0	DHS 2007
Women with low BMI (< 18.5 kg/m², %)			-			-	-	-	-	-	-	-	-