

Nutrition Profile

loqo

DEMOGRAPHICS

Total population (000)	6,459	(2008)
Total under-five population (000)	947	(2008)
Total number of births (000)	213	(2008)
Under-five mortality rate (per 1,000 live births)	98	(2008)
Total number of under-five deaths (000)	20	(2008)
Infant mortality rate (per 1,000 live births)	64	(2008)
Neonatal mortality rate (per 1,000 live births)	39	(2004)
HIV prevalence rate (15-49 years, %)	3.3	(2007)
Population below international poverty line of US\$1.25 per day (%)	39	(2006)






NUTRITIONAL STATUS



INFANT AND YOUNG CHILD FEEDING



Infant feeding practices, by age

48

2008

Other NS


28

2006

MICS

Togo

MICRONUTRIENTS




ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE

Pregnancy		Birth		0-5 months		6-23 months	>	24-59 months	
Use of iron-folic acid Early initiation of breastfeeding		53%	International Code of Marketing of Breastmilk Substitutes			Partial			
supplements	pplements (within 1 hour of birth)			Maternity protection in acc	ordanc	e with ILO Convention 183	No		
Household consumption of adequately iodized salt	25%	Infants not weighed at birth	56%	Exclusive breastfeeding (< 6 months)	- 48%				
			Continued breastfeeding at two years						
To increase children's chances of survival, improve development and prevent stunting, nutrition interventions need to be delivered during the mother's pregnancy and the					Full coverage of vitamin A supplementat		64%		
					National guidelines for management of severe acute malnutrition incorporating the community-based approach			Yes	
first two years of the child's life.						Policy on new ORS formula and zinc for management of diarrhoea*			
			ed	Policy on community treatment of pneumonia with antibiotics*					


MATERNAL NUTRITION/HEALTH

WATER AND SANITATION

510	(2005)
480	(1993-1998)
1,200	(2005)
38	(2005)
11	(1998)
38	(2008)
84	(2006)
46	(1998)
62	(2006)
12	(2006)
72, 82	(2007)
0.88	(2007)
	480 1,200 38 11 38 84 46 62 12 72, 82



Diarrhoea:



DISPARITIES IN NUTRITION

	Gender			Residence									
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	28	25	1.1		-	-			-	-	-	-	Other NS 2008
Underweight prevalence (WHO Child Growth Standards, %)	21	21	1.0	-	-				-				Other NS 2008
Wasting prevalence (WHO Child Growth Standards, %)	6	6	1.0		-	-	-		-	-	-		Other NS 2008
Infants not weighed at birth (%)			-	30	72	0.4	82	77	60	40	22	0.3	MICS 2006
Early initiation of breastfeeding (%)		-	-	40	34	1.2	29	29	43	40	40	1.4	MICS 2006
Women with low BMI ($<\!18.5kg/m^2\!,\%\!)$		11	-	9	12	0.8	-	-	-	-		-	DHS 1998

Under-five deaths caused by: