

Tonga

DEMOGRAPHICS

Total population (000) 104 (2008) Total under-five population (000) 14 (2008) Total number of births (000) 3 (2008) Under-five mortality rate (per 1,000 live births) 19 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 17 (2008) Neonatal mortality rate (per 1,000 live births) 12 (2004) HIV prevalence rate (15-49 years, %) . . Population below international poverty line of US\$1.25 per day (%) . .			
Total number of births (000) 3 (2008) Under-five mortality rate (per 1,000 live births) 19 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 17 (2008) Neonatal mortality rate (per 1,000 live births) 12 (2004) HIV prevalence rate (15-49 years, %)	Total population (000)	104	(2008)
Under-five mortality rate (per 1,000 live births) Total number of under-five deaths (000) Infant mortality rate (per 1,000 live births) Neonatal mortality rate (per 1,000 live births) HIV prevalence rate (15-49 years, %) Population below international	Total under-five population (000)	14	(2008)
births) 19 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 17 (2008) Neonatal mortality rate (per 1,000 live births) 12 (2004) HIV prevalence rate (15-49 years, %)	Total number of births (000)	3	(2008)
Infant mortality rate (per 1,000 live births) 17 (2008) Neonatal mortality rate (per 1,000 live births) 12 (2004) HIV prevalence rate (15-49 years, %) Population below international		19	(2008)
Neonatal mortality rate (per 1,000 live births) 12 (2004) HIV prevalence rate (15-49 years, %) Population below international	Total number of under-five deaths (000)	0	(2008)
births) 12 (2004) HIV prevalence rate (15-49 years, %) Population below international	Infant mortality rate (per 1,000 live births)	17	(2008)
Population below international		12	(2004)
·	HIV prevalence rate (15-49 years, %)		
	•	٠	





NUTRITIONAL STATUS

Burden of undernutrition (2008) WHO Child Growth Standards

Stunted (under-fives, 000):

Share of developing world stunting burden (%):

Stunting country rank:

- Underweight (under-fives, 000):

Wasted (under-fives, 000):

Severely wasted (under-fives, 000):

Current nutritional status

Percentage of children < 5 years old suffering

No Data

Stunting trends

Percentage of children < 5 years old stunted NCHS reference population

No Data

Underweight trends

Percentage of children < 5 years old underweight NCHS reference population


No Data

INFANT AND YOUNG CHILD FEEDING

Infant feeding practices, by age

No Data





Tonga

MICRONUTRIENTS


Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data

lodized salt consumption trends*
Percentage of households consuming adequately iodized salt

No Data





ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE




MATERNAL NUTRITION/HEALTH

WATER AND SANITATION







Under-five deaths caused by: Diarrhoea: 14% Pneumonia:

DISPARITIES IN NUTRITION

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)					-	•	-				-	•	
Underweight prevalence (WHO Child Growth Standards, %)	-	-									-		
Wasting prevalence (WHO Child Growth Standards, %)		-	•		-	-	-				-	•	
Infants not weighed at birth (%)	-	-			-						-	•	•
Early initiation of breastfeeding (%)	-	-	-		-		-	-			-		
Women with low BMI (< 18.5 kg/m², %)			-		-		-	-	-	-	-	-	