



# DEMOGRAPHICS

Total population (000)	234	(2008)
Total under-five population (000)	33	(2008)
Total number of births (000)	7	(2008)
Under-five mortality rate (per 1,000 live births)	33	(2008)
Total number of under-five	0	(2008)
Infant mortality rate (per 1,000 live births)	27	(2008)
Neonatal mortality rate (per 1,000 live births)	18	(2004)
HIV prevalence rate (15-49 years, %)		
Population below international poverty line of US\$1.25 per day (%)		








# NUTRITIONAL STATUS



Stunted (under-fives, 000):	7	Underweight (under-fives, 000):	5
Share of developing world stunting burden (%):	0.0	Wasted (under-fives, 000):	2
Stunting country rank:	129	Severely wasted (under-fives, 000):	0







### **INFANT AND YOUNG CHILD FEEDING**




# Vanuatu


#### **MICRONUTRIENTS**

#### Vitamin A supplementation

Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data






\* Estimates may not be comparable.


# **ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE**




## MATERNAL NUTRITION/HEALTH

# **WATER AND SANITATION**







Under-five deaths caused by:

Diarrhoea:

2% Pneumonia:

#### **DISPARITIES IN NUTRITION**

	Gender			Residence			Wealth quintile						
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-	-			-	-					-		
Underweight prevalence (WHO Child Growth Standards, %)		-									-		
Wasting prevalence (WHO Child Growth Standards, %)			•		-	•					-	•	•
Infants not weighed at birth (%)		-		9	23	0.4	34	20	22	9	9	0.3	MICS 2007
Early initiation of breastfeeding (%)				82	70	1.2	68	69	72	75	83	1.2	MICS 2007
Women with low BMI ( < 18.5 kg/m², %)		-	-					-	-	-	-	-	